arXiv:1101.1694v2 [math.OA] 13 Jan 2011

QUANTUM FUNCTIONS

ANDRE KORNELL

Abstract. Weaver has recently defined the notion of a quantum relation on a von Neumann algebra. We demonstrate that the corresponding notion of a quantum function between two von Neumann algebras coincides with that of a normal unital $*$-homomorphism in the opposite direction.

A relation between sets X and Y is simply a subset of $Y \times X$.[1] Weaver has recently proposed the following generalization of relations to the noncommutative setting:

Definition (Weaver, [5, Definition 2.1])**.** Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras. A *quantum relation* between $\mathcal{M}$ and $\mathcal{N}$ is an ultraweakly closed subspace $\mathcal{V} \subseteq \mathcal{B}(\mathcal{H}, \mathcal{K})$ such that $\mathcal{N}'\mathcal{V}\mathcal{M}' \subseteq \mathcal{V}$.

This definition reduces to the usual one when $\mathcal{M} = \ell^\infty(X)$ and $\mathcal{N} = \ell^\infty(Y)$. It has many other virtues. It is simple to state and simple to handle, and many familiar properties of relations have natural analogs.

Definition (Weaver, [5, Definition 2.4])**.** Let $\mathcal{M}_0$, $\mathcal{M}_1$ and $\mathcal{M}_2$ be von Neumann algebras.

- The *diagonal quantum relation* on $\mathcal{M}_0$ is the quantum relation $\mathcal{M}_0'$ between $\mathcal{M}_0$ and $\mathcal{M}_0$.
- If $\mathcal{V}$ is a quantum relation between $\mathcal{M}_0$ and $\mathcal{M}_1$, then the *inverse*[2] of $\mathcal{V}$ is the quantum relation $\mathcal{V}^*$ between $\mathcal{M}_1$ and $\mathcal{M}_0$.
- If $\mathcal{V}_0$ is a quantum relation between $\mathcal{M}_0$ and $\mathcal{M}_1$, and $\mathcal{V}_1$ is a quantum relation between $\mathcal{M}_1$ and $\mathcal{M}_2$, then their *composition* $\mathcal{V}_1 \circ \mathcal{V}_0$ is the quantum relation between $\mathcal{M}_0$ and $\mathcal{M}_2$ defined by

$$\mathcal{V}_1 \circ \mathcal{V}_0 = \mathcal{V}_1\mathcal{V}_0 = \overline{\operatorname{span}\{v_1 v_0 \mid v_1 \in \mathcal{V}_1,\, v_0 \in \mathcal{V}_0\}}^{ultraweak}.$$

If we interpret inclusion between quantum relations as the proper generalization of inclusion between classical relations, we arrive immediately at the following definitions.

Definition (Weaver, [5, Definition 2.4])**.** Let $\mathcal{V}$ be a quantum relation on a von Neumann algebra $\mathcal{M}$. Then $\mathcal{V}$ is said to be

- *reflexive* in case $\mathcal{M}' \subseteq \mathcal{V}$,
- *symmetric* in case $\mathcal{V}^* = \mathcal{V}$,
- *antisymmetric* in case $\mathcal{V} \cap \mathcal{V}^* \subseteq \mathcal{M}'$, and

---

The research reported here was supported by National Science Foundation grant DMS-0753228.

[1] Here and elsewhere, I have reversed the usual order of the Cartesian product so that the composition of two functions coincides with their composition as relations.

[2] Weaver uses *adjoint* and *product* in place of *inverse* and *composition*. I prefer the classical terminology.

- *transitive* in case $\mathcal{V}\mathcal{V} \subseteq \mathcal{V}$.

Thus, Weaver has generalized a large class of mathematical objects including orderings, graphs, equivalence relations, etc.

In foundations, a function from a set X to a set Y is typically defined as a relation $F \subseteq Y \times X$ such that for each element $x \in X$, there is exactly one element $y \in Y$ such that $(y, x) \in F$. Denoting the diagonal relations on X and Y by Δ_X and Δ_Y respectively, we may restate this condition as a pair of inequalities:

1. $\Delta_X \subseteq F^* \circ F$
2. $F \circ F^* \subseteq \Delta_Y$

It is therefore natural to investigate the quantum relations that satisfy the analogs of these inequalities, i.e., quantum functions. In fact, we show that they correspond exactly to the normal unital $*$-homomorphisms:

Definition. Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras. A *quantum function* from $\mathcal{M}$ to $\mathcal{N}$ is a quantum relation $\mathcal{V}$ between $\mathcal{M}$ and $\mathcal{N}$ such that $\mathcal{M}' \subseteq \mathcal{V}^*\mathcal{V}$ and $\mathcal{V}\mathcal{V}^* \subseteq \mathcal{N}'$.

Theorem. *Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras. There is a canonical bijective correspondence between normal unital $*$-homomorphisms $\mathcal{N} \longrightarrow \mathcal{M}$, and quantum functions from $\mathcal{M}$ to $\mathcal{N}$. This correspondence is functorial.*

Conventions used in this article. Let $\mathcal{H}$ be a Hilbert space. If $\xi \in \mathcal{H}$, then $\hat{\xi} \in \mathcal{B}(\mathbb{C}, \mathcal{H})$ is defined by $\hat{\xi}(c) = c\xi$. If $\mathcal{V}$ and $\mathcal{W}$ are ultraweakly closed subspaces of $\mathcal{B}(\mathcal{H})$, then $\mathcal{V}\mathcal{W} = \overline{\operatorname{span}\{vw \mid v \in \mathcal{V},\, w \in \mathcal{W}\}}^{uw}$, and $\mathcal{V}\bar{\otimes}\mathcal{W} = \overline{\operatorname{span}\{v \otimes w \mid v \in \mathcal{V},\, w \in \mathcal{W}\}}^{uw}$. The tensor product of two Hilbert spaces is defined in such a way that $\mathcal{H} \otimes \mathbb{C} = \mathcal{H} = \mathbb{C} \otimes \mathcal{H}$. The von Neumann algebra of scalar operators on $\mathcal{H}$ is denoted by $\mathbb{C}_{\mathcal{H}}$.

Definition. Let $\mathcal{M}$ and $\mathcal{N}$ be von Neumann algebras. Then $\mathbf{vN}(\mathcal{N}, \mathcal{M})$ denotes the set of normal unital $*$-homomorphisms $\mathcal{N} \longrightarrow \mathcal{M}$, and $\mathbf{qF}(\mathcal{M}, \mathcal{N})$ denotes the set of quantum functions from $\mathcal{M}$ to $\mathcal{N}$.

1. Functions from Homomorphisms

Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras, and let $\pi : \mathcal{N} \longrightarrow \mathcal{M}$ be a normal unital $*$-homomorphism.

Theorem 1.1 (Dixmier, [4, Theorem IV.5.5])**.** *There is a Hilbert space $\mathcal{L}$ and an isometry $w \in \mathcal{B}(\mathcal{H}, \mathcal{K} \otimes \mathcal{L})$ such that $\pi(b) = w^*(b \otimes 1)w$.*

Let $\mathcal{L}$ and w be as in Theorem 1.1 above.

Proposition 1.2. *For all $b \in \mathcal{N}$, $(b \otimes 1)w = w\pi(b)$.*

Proof. For all $b \in \mathcal{N}$,

$$ww^*(b^* \otimes 1)(b \otimes 1)ww^* = w\pi(b^*b)w^* = w\pi(b^*)\pi(b)w^* = ww^*(b^* \otimes 1)ww^*(b \otimes 1)ww^*,$$

so $ww^*(b^* \otimes 1)(1 - ww^*)(b \otimes 1)ww^* = 0$. Since $1 - ww^*$ is a projection, we conclude that $(1 - ww^*)(\mathcal{N}\bar{\otimes}\mathbb{C}_{\mathcal{L}})ww^* = 0$, i.e., $ww^* \in (\mathcal{N}\bar{\otimes}\mathbb{C}_{\mathcal{L}})'$. Therefore, for all $b \in \mathcal{N}$, $(b \otimes 1)w = (b \otimes 1)ww^*w = ww^*(b \otimes 1)w = w\pi(b)$. $\square$

Definition 1.3. The set $\mathfrak{G}(\pi) = \{v \in \mathcal{B}(\mathcal{H}, \mathcal{K}) \mid \forall b \in \mathcal{N}\ bv = v\pi(b)\}$ is a quantum relation between $\mathcal{M}$ and $\mathcal{N}$.

Proposition 1.4. $\mathcal{M}' \subseteq \mathfrak{G}(\pi)^*\mathfrak{G}(\pi)$

Proof. Choose a basis $\{e_\alpha\}_{\alpha\in I}$ of $\mathcal{L}$. For each $\alpha \in I$, $(1\otimes \hat{e}_\alpha^*)w \in \mathcal{B}(\mathcal{H},\mathcal{K})$ satisfies $b(1\otimes \hat{e}_\alpha^*)w = (1\otimes \hat{e}_\alpha^*)(b\otimes 1)w = (1\otimes \hat{e}_\alpha^*)w\pi(b)$ for all $b \in \mathcal{N}$, i.e., $(1\otimes \hat{e}_\alpha^*)w \in \mathfrak{G}(\pi)$. It follows that

$$1 = w^*w = \sum_{\alpha\in I} w^*(1\otimes \hat{e}_\alpha\hat{e}_\alpha^*)w = \sum_{\alpha\in I} \left((1\otimes \hat{e}_\alpha^*)w\right)^* \left((1\otimes \hat{e}_\alpha^*)w\right),$$

so $\mathbb{C}_\mathcal{H} \subseteq \mathfrak{G}(\pi)^*\mathfrak{G}(\pi)$. We conclude that $\mathcal{M}' = \mathbb{C}_\mathcal{H}\mathcal{M}' \subseteq \mathfrak{G}(\pi)^*\mathfrak{G}(\pi)\mathcal{M}' = \mathfrak{G}(\pi)^*\mathfrak{G}(\pi)$. ☐

Proposition 1.5. $\mathfrak{G}(\pi)\mathfrak{G}(\pi)^* \subseteq \mathcal{N}'$

Proof. For all $v_0, v_1 \in \mathfrak{G}(\pi)$, and all $b \in \mathcal{N}$, $bv_0v_1^* = v_0\pi(b)v_1^* = v_0(v_1\pi(b^*))^* = v_0v_1^*b$. ☐

Proposition 1.6. *Therefore* $\mathfrak{G}(\pi)$ *is a quantum function from* $\mathcal{M}$ *to* $\mathcal{N}$.

Thus, we have defined a function $\mathfrak{G} : \mathbf{vN}(\mathcal{N},\mathcal{M}) \longrightarrow \mathbf{qF}(\mathcal{M},\mathcal{N})$.

2. Homomorphisms from Functions

Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras, and $\mathcal{V}$ a quantum function from $\mathcal{M}$ to $\mathcal{N}$.

Lemma 2.1. *There exists a family*[3] $\{u_\alpha\}_{\alpha\in I}$ *of partial isometries in* $\mathcal{V}$ *such that*

1. *for all distinct* $\alpha, \beta \in I$, $u_\alpha u_\beta^* = 0$, *and*
2. $\sum u_\alpha^* u_\alpha = 1$.

Proof. Let $\mathcal{F}$ be the collection of all sets S of partial isometries in $\mathcal{V}$ such that $u\tilde{u}^* = 0$ whenever $u, \tilde{u} \in S$ are distinct. Applying Zorn's Lemma, we obtain a maximal such set $\hat{S}$.

Suppose that $\sum_{u\in\hat{S}} u^*u \neq 1$, and let $p = 1 - \sum_{u\in\hat{S}} u^*u$. The subspace $\mathcal{V}p$ is non-zero because $1 \in \mathcal{M}' \subseteq \mathcal{V}^*\mathcal{V}$. Therefore, pick $v \neq 0$ in $\mathcal{V}p \subseteq \mathcal{V}$. We will now obtain a partial isometry to add to $\hat{S}$ from the polar decomposition of v.

Let $W^*(\mathcal{V})$ be the von Neumann algebra generated by $\mathcal{V}$, which is the ultraweakly closed subspace of $\mathcal{B}(\mathcal{H}\oplus\mathcal{K})$ generated by finite words in the elements of $\mathcal{V}$ and their conjugates. Since $\mathcal{V}\mathcal{V}^*\mathcal{V} \subseteq \mathcal{N}'\mathcal{V} \subseteq \mathcal{V}$, $W^*(\mathcal{V})$ is in fact generated by words of one of the following forms: 1, v_0, v_0^*, $v_1^*v_0$, and $v_1v_0^*$. We conclude that $[\mathcal{K}]W^*(\mathcal{V})[\mathcal{H}] = \mathcal{V}$, where $[\mathcal{H}]$ and $[\mathcal{K}]$ denote projections onto $\mathcal{H}$ and $\mathcal{K}$ respectively.

Let $v = u_v|v|$ be the polar decomposition of x. Since $[\mathcal{K}]v[\mathcal{H}] = v$, $[\mathcal{K}]u_v[\mathcal{H}] = u_v$, so $u_v \in \mathcal{V}$. Furthermore, since $vp = v$, $u_vp = u_v$, so $u_vu^* = 0$ for all $u \in \hat{S}$. Thus, $\hat{S} \cup \{u_v\}$ is an element of $\mathcal{F}$ strictly larger than $\hat{S}$, a contradiction. ☐

Definition 2.2. Let $\mathfrak{G}^{-1}(\mathcal{V})$ be the normal unital $*$-homomorphism defined by

$$\mathfrak{G}^{-1}(\mathcal{V})(b) = \sum_{\alpha\in I} u_\alpha^* b u_\alpha = w_I^*(b\otimes 1)w_I,$$

where $\{u_\alpha\}_{\alpha\in I}$ is any family of partial isometries in $\mathcal{V}$ such that $\sum u_\alpha^*u_\alpha = 1$ and $u_\alpha u_\beta^* = 0$ whenever $\alpha \neq \beta$, and the isometry $w_I \in B(\mathcal{H}, \mathcal{K}\otimes\ell^2(I))$ is defined by $w_I\xi = \sum_{\alpha\in I} u_\alpha\xi \otimes e_\alpha$.

---

[3] Lemma 2.1 is a special case of Paschke's structural theorem for self-dual Hilbert W^*-modules ([3, Theorem 3.12]).

Proposition 2.3. *The normal unital $*$-homomorphism $\mathfrak{G}^{-1}(\mathcal{V})$ is well defined.*

Proof. Apply Lemma 2.1 above to obtain a family $\{u_\alpha\}_{\alpha\in I}$ of partial isometries in $\mathcal{V}$ such that $\sum u_\alpha^* u_\alpha = 1$ and $u_\alpha u_\beta^* = 0$ whenever $\alpha \neq \beta$. Let $w_I = \sum u_\alpha \otimes \hat{e}_\alpha \in \mathcal{B}(\mathcal{H}, \mathcal{K}\otimes \ell^2(I))$. For all $\xi \in \mathcal{H}$,

$$\|w_I(\xi)\|^2 = \sum_{\alpha,\beta} \langle u_\alpha(\xi)\otimes e_\alpha | u_\beta(\xi)\otimes e_\beta\rangle = \sum_\alpha \langle u_\alpha \xi | u_\alpha \xi\rangle = \langle \xi | \left(\sum_\alpha u_\alpha^* u_\alpha\right) \xi\rangle = \|\xi\|^2.$$

Thus, w_I is an isometry, and we may now define a normal unital completely positive map $\pi_I : \mathcal{N} \longrightarrow \mathcal{B}(\mathcal{H})$ by $\pi_I(b) = w_I^*(b\otimes 1)w_I = \sum u_\alpha^* b u_\alpha$. For all $b_0, b_1 \in \mathcal{N}$,

$$\begin{aligned}\pi_I(b_0)\pi_I(b_1) &= \left(\sum_\alpha u_\alpha^* b_0 u_\alpha\right)\left(\sum_\beta u_\beta^* b_1 u_\beta\right)\\ &= \sum_\alpha u_\alpha^* b_0 u_\alpha u_\alpha^* b_1 u_\alpha\\ &= \sum_\alpha u_\alpha^* b_0 b_1 u_\alpha u_\alpha^* u_\alpha\\ &= \sum_\alpha u_\alpha^* b_0 b_1 u_\alpha = \pi_I(b_0 b_1)\end{aligned}$$

because $u_\alpha u_\alpha^* \in \mathcal{V}\mathcal{V}^* \subseteq \mathcal{N}'$. We conclude that π_I is a normal unital $*$-homomorphism. For all $b \in \mathcal{N}$, $c \in \mathcal{M}'$,

$$c\pi_I(b) = \sum_{\alpha,\beta\in I} u_\alpha^* u_\alpha c u_\beta^* b u_\beta = \sum_{\alpha,\beta\in I} u_\alpha^* b u_\alpha c u_\beta^* u_\beta = \pi_I(b)c$$

because $u_\alpha c u_\beta^* \in \mathcal{V}\mathcal{M}'\mathcal{V}^* \subseteq \mathcal{V}\mathcal{V}^* \subseteq \mathcal{N}'$. By the Double Commutant Theorem, $\pi_I(\mathcal{N}) \subseteq \mathcal{M}$, so π_I may be viewed as a normal unital $*$-homomorphism $\mathcal{N} \longrightarrow \mathcal{M}$.

If $\{u_\alpha\}_{\alpha\in J}$ is another family of partial isometries that satisfies $\sum_{\alpha\in J} u_\alpha^* u_\alpha = 1$ and $u_\alpha u_\beta^* = 0$ for distinct $\alpha, \beta \in J$, we may obtain in the same way a normal unital $*$-homomorphism $\pi_J : \mathcal{N} \longrightarrow \mathcal{M}$. However, for all $b \in \mathcal{N}$,

$$\pi_J(b) = \sum_{\alpha\in I,\beta\in J} u_\alpha^* u_\alpha u_\beta^* b u_\beta = \sum_{\alpha\in I,\beta\in J} u_\alpha^* b u_\alpha u_\beta^* u_\beta = \pi_I(b)$$

because $u_\alpha u_\beta^* \in \mathcal{V}\mathcal{V}^* \subseteq \mathcal{N}'$. Thus, π_I is independent of our choice of family $\{u_\alpha\}_{\alpha\in I}$, and we may define $\mathfrak{G}^{-1}(\mathcal{V}) = \pi_I$. □

Thus, we have defined a function $\mathfrak{G}^{-1} : \mathbf{qF}(\mathcal{M},\mathcal{N}) \longrightarrow \mathbf{vN}(\mathcal{N},\mathcal{M})$.

3. $\mathfrak{G}^{-1}$ IS THE INVERSE OF $\mathfrak{G}$

Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras.

Proposition 3.1. *Let $\pi : \mathcal{N} \longrightarrow \mathcal{M}$ be a normal unital $*$-homomorphism. Then $\mathfrak{G}^{-1}(\mathfrak{G}(\pi)) = \pi$.*

Proof. Let $\{u_\alpha\}_{\alpha\in I}$ be a family of partial isometries in $\mathfrak{G}(\pi)$ such that $\sum_{\alpha\in I} u_\alpha^* u_\alpha = 1$ and $u_\alpha u_\beta^* = 0$ whenever $\alpha \neq \beta$. For all $b \in \mathcal{N}$,

$$\mathfrak{G}^{-1}(\mathfrak{G}(\pi))(b) = \sum_{\alpha\in I} u_\alpha^* b u_\alpha = \sum_{\alpha\in I} u_\alpha^* u_\alpha \pi(b) = \pi(b).$$

$\square$

Proposition 3.2. *Let* $\mathcal{L}$ *be a Hilbert space, and let* $w_0, w_1 \in \mathcal{B}(\mathcal{H}, \mathcal{K} \otimes \mathcal{L})$ *be isometries such that* $\pi_i(b) = w_i^*(b \otimes 1)w_i$ *defines a pair of* $*$*-homomorphisms* $\mathcal{N} \longrightarrow \mathcal{M}$*. If* $\pi_0 = \pi_1$*, then* $w_0 w_1^* \in (\mathcal{N} \overline{\otimes} \mathbb{C}_{\mathcal{L}})'$*.*

Proof. For all $b \in \mathcal{N}$, $(b \otimes 1)w_0 w_1^* = w_0 \pi_0(b) w_1^* = w_0 \pi_1(b) w_1^* = w_0(w_1 \pi_1(b^*))^* = w_0((b^* \otimes 1)w_1))^* = w_0 w_1^*(b \otimes 1)$. $\square$

Lemma 3.3. *Let* $\mathcal{V}$ *be a quantum function from* $\mathcal{M}$ *to* $\mathcal{N}$*, and let* $\{u_\alpha\}_{u \in I}$ *and* w_I *be as in Definition 2.2. Then* $(\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I)})' w_I \mathcal{M}' = \mathcal{V} \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I))$*.*

Proof. By definition, $w_I = \sum u_\alpha \otimes \hat{e}_\alpha \in \mathcal{V} \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I))$, so $(\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I)})' w_I \mathcal{M}' = (\mathcal{N}' \overline{\otimes} \mathcal{B}(\ell^2(I))) w_I \mathcal{M}' \subseteq \mathcal{V} \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I))$.

Let $f \in \ell^2(I)$ and $v \in \mathcal{V}$. Then for all $\alpha \in I$,

$$u_\alpha \otimes \hat{f} = \sum_{\beta \in I} (1 \otimes \hat{f} \hat{e}_\alpha^*)(u_\beta \otimes \hat{e}_\beta) = (1 \otimes \hat{f} \hat{e}_\alpha^*) w_I \in (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I)})' w_I \mathcal{M}',$$

so

$$v \otimes \hat{f} = \sum_{\alpha \in I} (v \otimes \hat{f})(u_\alpha^* u_\alpha) = \sum_{\alpha \in I} (v u_\alpha^* \otimes 1)(u_\alpha \otimes \hat{f}) \in (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I)})' w_I \mathcal{M}'$$

because $v u_\alpha^* \in \mathcal{V}\mathcal{V}^* \subseteq \mathcal{N}'$. It follows that $\mathcal{V} \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I)) \subseteq (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I)})' w_I \mathcal{M}'$, concluding the proof. $\square$

Proposition 3.4. *The function* $\mathfrak{G}^{-1}$ *is injective.*

Proof. For $k \in \{0, 1\}$, let $\mathcal{V}_k \in \mathbf{qF}(\mathcal{M}, \mathcal{N})$, and let $\{u_\alpha\}_{\alpha \in I_k}$ be a family of partial isometries in $\mathcal{V}_k$ such that $u_\alpha u_\beta^* = 0$ for distinct $\alpha, \beta \in I_k$, and $\sum u_\alpha^* u_\alpha = 1$. We may assume that I_0 and I_1 have equal, non-zero cardinality by throwing in indexed instances of the zero partial isometry where necessary. Thus, we may choose a unitary $s \in \mathcal{B}(\ell^2(I_0), \ell^2(I_1))$.

Suppose that $\mathfrak{G}^{-1}(\mathcal{V}_0) = \mathfrak{G}^{-1}(\mathcal{V}_1)$. For all $b \in \mathcal{N}$,

$$\begin{aligned} ((1 \otimes s) w_{I_0})^* (b \otimes 1)((1 \otimes s) w_{I_0}) = w_{I_0}^* (b \otimes 1) w_{I_0} &= \mathfrak{G}^{-1}(\mathcal{V}_0)(b) \\ &= \mathfrak{G}^{-1}(\mathcal{V}_1)(b) = w_{I_1}^* (b \otimes 1) w_{I_1} \end{aligned}$$

By Proposition 3.2, $(1 \otimes s) w_{I_0} w_{I_1}^* \in (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I_1)})'$.

By Lemma 3.3 above,

$$\begin{aligned} \mathcal{V}_1 \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I_1)) &= (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I_1)})' w_{I_1} \mathcal{M}' \\ &\supseteq (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I_1)})' ((1 \otimes s) w_{I_0} w_{I_1}^*) w_{I_1} \mathcal{M}' \\ &= (\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I_1)})' (1 \otimes s) w_{I_0} \mathcal{M}' \\ &= (1 \otimes s)(\mathcal{N} \overline{\otimes} \mathbb{C}_{\ell^2(I_0)})' w_{I_0} \mathcal{M}' \\ &= (1 \otimes s)(\mathcal{V}_0 \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I_0))) \\ &= \mathcal{V}_0 \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I_1)). \end{aligned}$$

Choosing an arbitrary unit vector $f \in \ell^2(I_1)$, we conclude that

$$\mathcal{V}_1 = (1 \otimes \hat{f}^*) \left(\mathcal{V}_1 \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I_1)) \right) \supseteq (1 \otimes \hat{f}^*) \left(\mathcal{V}_0 \overline{\otimes} \mathcal{B}(\mathbb{C}, \ell^2(I_1)) \right) = \mathcal{V}_0.$$

Similarly, $\mathcal{V}_1 \subseteq \mathcal{V}_0$, so $\mathcal{V}_1 = \mathcal{V}_0$. $\square$

Theorem 3.5. *Let $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ and $\mathcal{N} \subseteq \mathcal{B}(\mathcal{K})$ be von Neumann algebras. The function*

$$\mathfrak{G} : \mathbf{vN}(\mathcal{N}, \mathcal{M}) \longrightarrow \mathbf{qF}(\mathcal{M}, \mathcal{N})$$

defined by

$$\mathfrak{G}(\pi) = \{v \in \mathcal{B}(\mathcal{H}, \mathcal{K}) \,|\, \forall b \in \mathcal{N}\ bv = \pi(b)v\}$$

is a bijection.

Proof. The theorem follows immediately from Propositions 3.1 and 3.4. □

4. Functoriality

Proposition 4.1. *Let $\mathcal{M}_0$, $\mathcal{M}_1$, and $\mathcal{M}_2$ be von Neumann algebras. If $\mathcal{V}_0 \in \mathbf{qF}(\mathcal{M}_0, \mathcal{M}_1)$ and $\mathcal{V}_1 \in \mathbf{qF}(\mathcal{M}_1, \mathcal{M}_2)$, then $\mathcal{V}_1\mathcal{V}_0 \in \mathbf{qF}(\mathcal{M}_0, \mathcal{M}_2)$.*

Proof.

$$\mathcal{M}_0' \subseteq \mathcal{V}_0^*\mathcal{V}_0 \subseteq \mathcal{V}_0^*\mathcal{M}_1'\mathcal{V}_0 \subseteq \mathcal{V}_0^*\mathcal{V}_1^*\mathcal{V}_1\mathcal{V}_0 = (\mathcal{V}_1\mathcal{V}_0)^*(\mathcal{V}_1\mathcal{V}_0)$$
$$(\mathcal{V}_1\mathcal{V}_0)(\mathcal{V}_1\mathcal{V}_0)^* = \mathcal{V}_1\mathcal{V}_0\mathcal{V}_0^*\mathcal{V}_1 \subseteq \mathcal{V}_1\mathcal{M}_1'\mathcal{V}_1^* \subseteq \mathcal{V}_1\mathcal{V}_1^* \subseteq \mathcal{M}_2'$$

□

Proposition 4.2. *Let $\mathcal{M}_0$, $\mathcal{M}_1$, and $\mathcal{M}_2$ be von Neumann algebras. If $\pi_1 : \mathcal{M}_2 \longrightarrow \mathcal{M}_1$ and $\pi_0 : \mathcal{M}_1 \longrightarrow \mathcal{M}_0$ are normal unital $*$-homomorphisms, then $\mathfrak{G}(\pi_0 \circ \pi_1) = \mathfrak{G}(\pi_1)\mathfrak{G}(\pi_0)$.*

Proof. Clearly, $\mathfrak{G}(\pi_1)\mathfrak{G}(\pi_0) \subseteq \mathfrak{G}(\pi_0 \circ \pi_1)$. By Definition 2.2, $\mathfrak{G}^{-1}(\mathfrak{G}(\pi_1)\mathfrak{G}(\pi_0)) = \mathfrak{G}^{-1}(\mathfrak{G}(\pi_0 \circ \pi_1))$. By Theorem 3.5, we conclude that $\mathfrak{G}(\pi_1)\mathfrak{G}(\pi_0) = \mathfrak{G}(\pi_0 \circ \pi_1)$. □

Proposition 4.3. *Let $\mathcal{M}$ be a von Neumann algebra, and let $\iota : \mathcal{M} \longrightarrow \mathcal{M}$ be the identity $*$-homomorphism. Then $\mathfrak{G}(\iota) = \mathcal{M}'$.*

Proof. This is an immediate consequence of Definition 1.3. □

Definition 4.4. Let **vN** be the category whose objects are von Neumann algebras, and whose morphisms are normal unital $*$-homomorphisms.

Definition 4.5. Let **qF** be the following category:

- The objects of **qF** are von Neumann algebras.
- For any two objects $\mathcal{M}$ and $\mathcal{N}$, a morphism from $\mathcal{M}$ to $\mathcal{N}$ is a quantum function from $\mathcal{M}$ to $\mathcal{N}$.
- For any two morphisms $\mathcal{V}_0 \in \mathbf{qF}(\mathcal{M}_0, \mathcal{M}_1)$ and $\mathcal{V}_1 \in \mathbf{qF}(\mathcal{M}_1, \mathcal{M}_2)$, $\mathcal{V}_1 \circ \mathcal{V}_0 = \mathcal{V}_1\mathcal{V}_0$.
- For any object $\mathcal{M}$, the identity morphism at $\mathcal{M}$ is the quantum function $\mathcal{M}' \in \mathbf{qF}(\mathcal{M}, \mathcal{M})$.

Theorem 4.6. *The functor $\mathfrak{G} : \mathbf{vN} \longrightarrow \mathbf{qF}$ defined by*

- *for all objects $\mathcal{M}$ of* **vN**, *$\mathfrak{G}(\mathcal{M}) = \mathcal{M}$, and*
- *for all morphisms $\pi \in \mathbf{vN}(\mathcal{N} \subseteq \mathcal{B}(\mathcal{K}), \mathcal{M} \subseteq \mathcal{B}(\mathcal{H}))$,*

$$\mathfrak{G}(\pi) = \{v \in \mathcal{B}(\mathcal{H}, \mathcal{K}) \,|\, \forall b \in \mathcal{N}\ bv = v\pi(b)\},$$

is a coisomorphism of categories.

Proof. This is a straightforward consequence of Proposition 4.2, Proposition 4.3, and Theorem 3.5. □

References

[1] G. Kuperberg and N. Weaver, *A Von Neumann Algebra Approach to Quantum Metrics*, arXiv:1005.0353v2.

[2] V. M. Manuilov and E. V. Troitsky, *Hilbert C^*-Modules*, Translations of Mathematical Monographs **226** (2005).

[3] W. L. Paschke, *Inner Product Modules over B^*-algebras*, Transactions of the Americal Mathematical Society **182** (1973), 443-468.

[4] M. Takesaki, *Theory of Operator Algebras I*, Springer, 1979.

[5] N. Weaver, *Quantum Relations*, arXiv:1005.0354v1.

Department of Mathematics, University of California, Berkeley, CA 94720-3840
E-mail address: `kornell@math.berkeley.edu`